

PCP Constipation Referral Checklist

Patient Name _____
Date of Birth _____

Red Flag(s) present? (check all applicable answers below)

- | | |
|--|---|
| <input type="checkbox"/> Age < 1 month | <input type="checkbox"/> Perianal fistula |
| <input type="checkbox"/> Meconium > 48 hours | <input type="checkbox"/> Abnormal position of anus |
| <input type="checkbox"/> Family history of Hirschsprung's | <input type="checkbox"/> Absent anal or cremasteric reflex |
| <input type="checkbox"/> Ribbon stools | <input type="checkbox"/> Decreased lower extremity strength/tone/reflex |
| <input type="checkbox"/> Blood in the stools in absence of anal fissures | <input type="checkbox"/> Tuft of hair on spine |
| <input type="checkbox"/> Failure to thrive | <input type="checkbox"/> Sacral dimple |
| <input type="checkbox"/> Biliious vomiting | <input type="checkbox"/> Gluteal cleft deviation |
| <input type="checkbox"/> Abnormal thyroid gland | <input type="checkbox"/> Extreme fear during anal inspection |
| <input type="checkbox"/> Severe abdominal distension | <input type="checkbox"/> Anal scars |

Osmotic AND Stimulant laxatives used for ≥ 1 month? (check all that apply)

Osmotic laxatives: Miralax Lactulose/Kristalose Milk of Mag

Stimulant laxatives: Exlax Senna Dulcolax

Other: _____

Clean out attempted? If yes, check indication below


- soiling (encopresis) impaction suspected
 colonic reset needed

Other: _____

Clean Out Regimen Options (*Goal: transparent liquid stools without sediment*)

- | | | |
|------------------|-----------------------------|---|
| Option 1: | <i>Enemas Only:</i> | <u>Age < 2 y/o:</u> Peds Fleets enema daily x 3 days; <u>Age > 2 y/o:</u> Adult fleets enema daily x 3 days; may need 1-2 additional days |
| Option 2: | <i>Enemas + Miralax:</i> | Mineral oil enema, Fleets enema followed by Miralax q 30 min to 1 hour for 4 hours |
| Option 3: | <i>Oral laxatives only:</i> | 8 am to 8 pm, clear liquid diet (eat BF before 8 am, dinner after 8 pm)
<u>0800 and 2000:</u> dulcolax/senna + miralax; <u>1000, noon, 1400, 1600, 1800:</u> miralax only (*max of 2 consecutive days) |
| Option 4: | <i>Oral laxatives only:</i> | Senna + Miralax + Senna (over 2 hours). 1 day only |

Algorithm for Primary Care Providers



Laxative Dosing Guideline

Oral laxatives

Dosages

Osmotic laxatives

Lactulose

1–2 g/kg, once or twice/day

Maintenance: $0.2\text{--}0.8 \text{ g} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$

Fecal disimpaction: $1\text{--}1.5 \text{ g} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ (with a maximum of 6 consecutive days)

2–5 y: 0.4–1.2 g/day, once or divided

6–11 y: 1.2–2.4 g/day, once or divided

12–18 y: 2.4–4.8 g/day, once or divided

Miralax (PEG or polyethylene glycol)

Milk of magnesia (magnesium hydroxide)

Fecal softeners

Mineral oil

1–18 y: $1\text{--}3 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$, once or divided, max 90 mL/day

Stimulant laxatives

Bisacodyl

3–10 y: 5 mg/day

>10 y: 5–10 mg/day

Senna

12 mo–6 y: 2.5–5 mg once or twice/day

6–12 y: 7.5–10 mg/day

>12 y: 15–20 mg/day

Sodium picosulfate

1 mo–4 y: 2.5–10 mg once/day

4–18 y: 2.5–20 mg once/day

Rectal laxatives/enemas

Bisacodyl

2–10 y: 5 mg once/day

>10 y: 5–10 mg once/day

Sodium docusate

<6 y: 60 mL

>6 y: 120 mL

Sodium phosphate

1–18 y: 2.5 mL/kg, max 133 mL/dose

NaCl

Neonate <1 kg: 5 mL, >1 kg: 10 mL

>1 y: 6 mL/kg once or twice/day

Mineral oil

2–11 y: 30–60 mL once/day

>11 y: 60–150 mL once/day